

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**

THIS PAGE BLANK (USPTO)



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 25/30, 47/36, 43/60, 43/40, 43/42	A1	(11) International Publication Number: WO 95/18531 (43) International Publication Date: 13 July 1995 (13.07.95)
(21) International Application Number: PCT/IB95/00001 (22) International Filing Date: 3 January 1995 (03.01.95) (30) Priority Data: 64/94-6 10 January 1994 (10.01.94) CH (71) Applicant (for all designated States except US): CIBA-GEIGY AG [CH/CH]; Klybeckstrasse 141, CH-4002 Basle (CH). (72) Inventors; and (75) Inventors/Applicants (for US only): VOGT, Manfred [DE/DE]; Rheinstrasse 25, D-79713 Bad Säckingen (DE). HUDETZ, Manfred [DE/CH]; Rheinweg 42, CH-4310 Rheinfelden (CH). (74) Common Representative: CIBA-GEIGY AG; Patentabteilung, Klybeckstrasse 141, CH-4002 Basle (CH).		(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MX, NO, NZ, PL, RO, RU, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: WETTABLE POWDER FORMULATIONS OF HERBICIDES (57) Abstract <p>A formulation comprising: (a) at least one herbicide, (b) at least one porous solid carrier material, and (c) a nonionic surfactant, which formulation contains 30 to 60 % by weight, based on the total formulation, of at least one nonionic surfactant of the formula (I): $R_1-(O-R_2)_n-O-R_3$, wherein R_1 is C_8-C_{16}alkyl or C_8-C_{16}alkenyl, R_2 is identical or different C_2-C_4 alkylene, R_3 is hydrogen or C_1-C_4alkyl, and n is an integer from 5-10. High biological activity is achieved with the solid formulation at low rates of application.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Larvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

Wettable powder formulations of herbicides

The present invention relates to a wettable powder formulation of solid herbicides on a solid carrier material and large amounts of a nonionic surfactant which is a reaction product of an alcohol with longer chain length and identical or different alkylene oxides, and whose end groups may additionally be alkylated.

Pesticides are usually marketed as formulated concentrates. In addition to water-dispersible powders, it is also possible to use suspension concentrates, emulsifiable concentrates or other physical forms, as described, inter alia, in EP-B-0 190 995. Emulsifiable concentrates consist essentially of a concentrated solution of the active ingredient in a suitable organic solvent, typically a mineral oil or xylene, which can be emulsified in water by adding suitable surfactants. They usually have very good biological activity and are easily to handle when formulating the spray mixture. One disadvantage is the use of an organic solvent.

Water-dispersible and wettable powders typically consist of inert carrier materials, one or more active ingredients, and surface active compounds that ensure easily wettability with water. Organic solvents are not required. It is frequently observed that these powders have a markedly lower biological activity than emulsifiable concentrates, so that this type of formulation is impracticable, i.e. it must be used in too high concentration.

Nonionic surfactants derived from fatty alcohol ethoxylates may also be used for formulating wettable powders. Usually, however, only minor amounts of surfactant are used, as the powders are required to be readily flowable and must remain storage-stable over a considerable period of time. On contact with water, no lumps shall form and the powder must form a fine dispersion as rapidly as possible when stirred in water.

C. L. McGiness, P. J. Porpiglia and G. R. Gillespie report in Abstr. Meet. Weed Sci. Soc. Am. (32, 13, 1992) on greater biological control when using primisulfuron in admixture with ethoxylated fatty alcohols. The fatty alcohol ethoxylate preferably consists of a 70 % ethoxylated C₁₂-C₁₄fatty alcohol. The fatty alcohol ethoxylate is mixed direct with the herbicide, the mixture is diluted with water and then applied in a greenhouse.

A wettable powder useful in practice having a high content of fatty alcohol ethoxylate and very good biological properties has so far not been disclosed.

- 2 -

It has now been found that storage-stable wettable powders of high biological activity can be obtained by applying herbicidally active compounds together with large amounts of an alkylene-oxylated alcohol with longer chain length as nonionic surfactant on a solid porous carrier material, even if said nonionic surfactant carries alkylated end groups.

Accordingly, in one of its aspects the invention relates to a formulation comprising

- a) at least one herbicide,
- b) at least one porous solid carrier material, and
- c) a nonionic surfactant,

which formulation contains 30 to 60 % by weight, based on the total formulation, of at least one nonionic surfactant of formula I



wherein R_1 is C_8 - C_{16} alkyl or C_8 - C_{16} alkenyl, R_2 is identical or different C_2 - C_4 alkylene, R_3 is hydrogen or C_1 - C_4 alkyl, and n is an integer from 5-10.

Suitable herbicides are known per se and described, inter alia, in R. Wegler, Chemie der Pflanzenschutz- und Schädlingsbekämpfungsmittel (Chemistry of Plant Protection Agents and Pesticides), Volume 2, Springer Verlag 1970, pp. 196-390.

Preferred herbicides are those selected from the groups consisting of the urea derivatives, imidazolinones, sulfonylureas, uracils, cyclohexanediones, chloroacetanilides, oxime ethers, triazines and phenoxypropionic acid derivatives.

Particularly preferred herbicides are those of the group of phenoxypropionic acid derivatives disclosed, inter alia, in EP 0 083 556. They are particularly useful active ingredients for the wettable powders of this invention.


Another preferred group of herbicides is that of the group of the sulfonylureas disclosed, inter alia, in EP-A-0 044 807 or in EP-A-0 120 814.

The novel formulations may additionally comprise safeners, typically those of the group of the quinoline derivatives. Particularly preferred safener-herbicide combinations are disclosed, inter alia, in EP-A-0 191 736 or in EP-A-0 094 349.


Particularly preferred herbicides are compounds of formula II which may be in the form of

- 3 -

the R enantiomer or the racemate



wherein Q is a radical of formula




wherein R_4 is hydrogen, F, Cl or Br;

R_5 is CF_3 , CN, Cl, Br or I;

R_7 and R_8 are each independently of the other F, Cl, Br, I, CF_3 or CN,

X is oxygen or sulfur, and n is 0 or an integer from 1 to 3;



wherein R_6 is hydrogen, $\text{C}_1\text{-C}_8\text{alkyl}$, $\text{C}_3\text{-C}_8\text{alkenyl}$, $\text{C}_3\text{-C}_8\text{alkynyl}$, $\text{C}_1\text{-C}_8\text{haloalkyl}$, $\text{C}_3\text{-C}_8\text{haloalkenyl}$, $\text{C}_3\text{-C}_8\text{haloalkynyl}$, or $\text{C}_1\text{-C}_6\text{alkyl}$ which is substituted by CN, NO_2 , $\text{C}_1\text{-C}_4\text{alkoxy}$, $\text{C}_1\text{-C}_4\text{alkyl-SO-}$, $\text{C}_1\text{-C}_4\text{alkyl-SO}_2\text{-}$, $\text{C}_1\text{-C}_8\text{alkoxycarbonyl}$, $\text{C}_3\text{-C}_8\text{alkenyloxycarbonyl}$, $\text{C}_3\text{-C}_8\text{alkynyloxycarbonyl}$, $\text{C}_1\text{-C}_4\text{alkylidene-aminoxy}$ or $\text{N(R}_{10}\text{)(R}_{11}\text{)}$, or is $\text{C}_3\text{-C}_8\text{alkenyl}$;

R_9 alkyl, haloalkyl, CN or $\text{C}_1\text{-C}_4\text{alkoxycarbonyl}$;

and R_{10} and R_{11} are each independently of the other $\text{C}_1\text{-C}_8\text{alkyl}$, $\text{C}_3\text{-C}_8\text{alkenyl}$,

- 4 -

C₃-C₈alkynyl, phenyl, benzyl or phenyl or benzyl, each substituted by halogen, NO₂, CN, OCH₃, or CF₃, or R₁₀ and R₁₁, together with the linking nitrogen atom, form a 5-7-membered saturated heterocycle.

Alkyl in R₆, R₉, R₁₀ and R₁₁ may be linear or branched and, in the significance of C₁-C₈alkyl, is typically ethyl, the different isomers of propyl, butyl, pentyl, hexyl, heptyl or octyl.



Alkenyl in R₆, R₁₀ and R₁₁ may be linear or branched and in the significance of C₃-C₈alkenyl is typically the different isomers of propenyl, butenyl, pentenyl, hexenyl, heptenyl or octenyl. Alkenyl is preferably C₂-C₇alkenyl-CH₂-.

Alkynyl may be linear or branched and in the significance of C₃-C₈alkynyl is typically the different isomers of propynyl, butynyl, pentynyl, hexynyl, heptynyl or octynyl. Alkynyl is preferably C₂-C₇alkynyl-CH₂-.

C₁-C₄Alkoxy is typically methoxy, ethoxy, propoxy or butoxy. C₁-C₄Alkylthio may be methylthio, ethylthio, propylthio or butylthio.

The 3-8-membered N-heterocycle is preferably morpholino or piperidino.




Very particularly preferred herbicides are compounds of formula III or IV



especially in the R-configuration.


Further preferred compounds are those of formula V, VI or VII.

- 5 -



Suitable safeners are a number of the quinoline derivatives disclosed in the references cited hereinabove. A compound of formula

- 6 -



(VIII)

is particularly suitable.

The herbicides can be used in an amount of 1 to 30 % by weight, preferably of 5 to 15 %, based on the entire formulation.

If a safener is concurrently used, the ratio of safener to herbicide is 1:5 to 5:1, preferably 1:4 to 1:1.

Herbicide and safener usually have an average particle size of 1:100, preferably of 1:50 and, most preferably, of 1 to 25 μm .

The solid carriers are porous and preferably have an inner surface area of at least 10 m^2/g , preferably 20 to 400 m^2/g and, most preferably, 20 to 200 m^2/g . Solid carriers for wettable powders are usually natural mineral fillers such as calcite, talcum, kaolin, diatomaceous earth, montmorillonite or attapulgite. In order to improve the physical properties it is also possible to add highly dispersed silicic acid, synthetic highly absorbent silicic acid, or porous polymers such as urea/formaldehyde condensate (Pergopak). Suitable granulated adsorptive carriers are porous types such as pumice, broken brick, sepiolite or bentonite; and suitable nonsorbent carriers are materials such as calcite or sand. In addition, a great number of pregranulated materials of inorganic or organic nature can be used, e.g. especially dolomite or pulverised plant residues. The carrier material usually has an average particle size of 1 to 200 μm , preferably 1 to 100 μm and, most preferably, 1 to 25 μm . The particle size of plant residues is usually larger and can be up to 500 μm .

The porous carrier material is preferably used in an amount of 30 to 60 % by weight, most preferably 35 to 50 % by weight, based on the total formulation. The percentages in connection with the formulation make up 100 %.

- 7 -

R_1 in formula I is preferably C_{10} - C_{15} alkyl and, most preferably, C_{12} - C_{14} alkyl. Alkyl may be linear or branched and is preferably linear. Alkyl may be mixtures of alkyl radicals with different chain lengths. Typical examples are octyl, nonyl, decyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl and hexadecyl.

n in formula I is preferably an integer from 6 to 8. The compounds of formula I are preferably technical mixtures that contain oligomers in which $n = 5-10$, preferably $n = 6-8$.

R_2 is typically ethylene, 1,2-propylene or 1,2-butylene. If R_2 denotes different alkylene radicals, the compounds may be random or block oligomers.

R_2 is typically a mixture of C_2 alkylene and C_4 alkylene and R_3 is hydrogen or methyl.

Most preferably R_1 is C_{10} - C_{15} alkyl, R_2 is C_2 alkylene, R_3 is hydrogen and n is an integer from 6 to 8.

The novel formulations may additionally comprise other surface-active compounds, conveniently in an amount of 1 to 15 % by weight, preferably 1 to 10 % by weight, based on the total formulation.

Depending on the type of active ingredient or combination of active ingredients to be formulated, suitable surface-active compounds are nonionic, cationic and/or anionic surfactants having good emulsifying, dispersing and wetting properties. Surfactants will also be understood as comprising mixtures of surfactants.

Suitable anionic surfactants may be water-soluble soaps as well as water-soluble synthetic surface-active compounds.

Suitable soaps are the alkali metal salts, alkaline earth metal salts, ammonium salts or substituted ammonium salts of higher fatty acids (C_{10} - C_{22}), e.g. the sodium or potassium salts of oleic or stearic acid, or of mixtures of natural fatty acids which can be obtained, inter alia, from coconut oil or tall oil. Further suitable soaps are also the fatty acid methyl taurin salts.

More often, however, so-called synthetic surfactants are used, especially fatty sulfonates, fatty sulfates, sulfonated benzimidazole derivatives or alkylarylsulfonates.

The fatty alcohol sulfonates or sulfates are usually in the form of alkali metal salts, alkaline earth metal salts, ammonium salts or substituted ammonium salts, and they contain a C₈-C₂₂alkyl radical which also includes the alkyl moiety of acyl radicals, e.g. the sodium or calcium salt of ligninsulfonic acid, of dodecylsulfate, or of a mixture of fatty alcohol sulfates obtained from natural fatty acids. These compounds also comprise the salts of sulfated and sulfonated fatty alcohol/ethylene oxide adducts. The sulfonated benzimidazole derivatives preferably contain two sulfonic acid groups and one fatty acid radical containing 8 to 22 carbon atoms. Illustrative examples of alkylarylsulfonates are the sodium, calcium or triethanolamine salts of dodecylbenzenesulfonic acid, dibutyl-naphthalenesulfonic acid, or of a condensate of naphthalenesulfonic acid and formaldehyde. Corresponding phosphates, typically salts of the phosphoric acid ester of an adduct of p-nonylphenol with 4 to 14 mol of ethylene oxide, or phospholipids, are also suitable.

Further suitable nonionic surfactants are preferably polyglycol ether derivatives of aliphatic or cycloaliphatic alcohols or of saturated or unsaturated fatty acids and alkylphenols, said derivatives containing 3 to 30 glycol ether groups and 8 to 20 carbon atoms in the (aliphatic) hydrocarbon moiety and 6 to 18 carbon atoms in the alkyl moiety of the alkylphenols. Further suitable nonionic surfactants are the water-soluble polyadducts of polyethylene oxide with polypropylene glycol, ethylenediaminopolypropylene glycol and alkylpolypropylene glycol containing 1 to 10 carbon atoms in the alkyl chain, which polyadducts contain 20 to 250 ethylene glycol ether groups and 10 to 100 propylene glycol ether groups. These compounds usually contain 1 to 5 ethylene glycol units per propylene glycol unit.

Illustrative examples of nonionic surfactants are nonylphenol polyethoxylates, polyethoxylated castor oil, polyadducts of polypropylene and polyethylene oxide, tributylphenol polyethoxylate, polyethylene glycol and octylphenol polyethoxylate. Fatty acid esters of polyoxyethylene sorbitan are also suitable nonionic surfactants, typically polyoxyethylene sorbitan trioleate.

Cationic surfactants are preferably quaternary ammonium salts carrying, as N-substituent, at least one C₈-C₂₂alkyl radical and, as further substituents, unsubstituted or halogenated lower alkyl, benzyl or hydroxy-lower alkyl radicals. The salts are preferably in the form of halides, methyl sulfates or ethyl sulfates, for example stearyl trimethylammonium chloride or benzyl bis(2-chloroethyl)ethylammonium bromide.

The surfactants customarily employed in the art of formulation are described, inter alia, in "Mc Cutcheon's Detergents and Emulsifiers Annual", MC Publishing Corp., Glen Rock, New Jersey, 1981, H. Stache, "Tensid-Taschenbuch" (Handbook of Surfactants), Carl Hanser Verlag, Munich/Vienna 1981, and M. and J. Ash. "Encyclopedia of Surfactants", Vol. I-III, Chemical Publishing Co., New York, 1980-1981.

The compositions are prepared in known manner, e.g. by homogeneously mixing and/or grinding the active ingredients with extenders, typically with solvents, solid carriers and, in some cases, surface-active compounds (surfactants).

Mixing can be carried out in ordinary commercial mixers. If the active ingredients have a low melting point, it is expedient to premix a portion of the carrier material with the total amount of herbicide and the safener, if used, and to grind the mixture, then to add the rest of the carrier material and, finally, to add the surfactant useful in the practice of this invention and other optional adjuvants.

It is usually often possible to mix the carrier loaded with the surfactant with the herbicide and afterwards to grind the mixture.

If the active ingredients have higher melting points, components (a) and (b) are conveniently mixed in the finely ground state or ground together and component (c) is then added.

The end user will normally use the active ingredient concentrates or wettable powders in the form of dilute compositions. This is done by dispersing a biologically active amount of the formulation in water.

The invention therefore also relates to a composition for controlling weeds and grasses in crops of cultivated plants, which composition contains a biologically effective amount of the novel formulation in water.

The amount of formulation added to water to prepare a spray mixture will depend on the type of application. Usually a 0.1 to 10 % by weight aqueous dispersion is prepared.

Suitable cultivated plants are preferably those which are important in the food or textile

- 10 -

sector, typically sugar cane and, in particular, millet, maize, rice and cereals (wheat, rye, barley, oats). The use of the novel composition in crops of cereals is very particularly preferred.

The weeds to be controlled can be both monocot as well as dicot weeds.

The concentration of herbicide will usually be from 0.001 to 1 kg/ha, but preferably from 0.01 to 0.5 kg a.i./ha. In postemergence treatment of the weeds, a reduction of 80 to 98 % of biomass is usually achieved.

Application of these compositions may be spraying, atomising, coating or drenching.

The invention also relates to a method of selectively controlling weeds, which comprises treating crops of cultivated plants, the seeds or the crop area thereof with an effective amount, usually from 0.01 to 0.5 kg/ha, of a herbicidal composition contains the herbicidal formulation of this invention, preferably using water as diluent.

The invention further relates to the use of nonionic surfactants of formula I for the preparation of wettable powder formulations for use as herbicidal compositions.

The invention is illustrated by the following Examples.

A) Formulation Examples

A premix comprising about 20 % by weight of herbicide and 5 % by weight of safener is prepared from the total amount of herbicide and safener in a mixing and milling apparatus with a portion of the carrier material. The remainder of the carrier material is then added and then the surfactant is applied. Afterwards the solid dispersants are added and a homogeneous mixture is prepared. The formulations are described in the following Examples. All percentages are by weight. The suspensibility of the wettable powders indicated in Examples A4 to A6 is determined in accordance with the method laid down in CIPAC MT 15.1.

Example A1 (formulation of a wettable powder)

8 % of compound of formula III

2 % of compound of formula VIII

- 11 -

3 % of sodium butylnaphthalenesulfonate

5 % of sodium ligninsulfonate

44 % of isotridecanol-8-ethoxylate (Gezetol 138[®])

38 % of highly dispersed urea-formaldehyde polycondensate (Pergopak[®] M)

Example A2 (formulation of a wettable powder)

4 % of compound of formula III

1 % of compound of formula VIII

1.5 % of sodium butylnaphthalenesulfonate

2.5 % sodium ligninsulfonate

55 % of isotridecanol-8-ethoxylate

36 % highly dispersed urea-formaldehyde polycondensate

Example A3 (formulation of a wettable powder)

8 % of compound of formula III

2 % of compound of formula VIII

2 % of epoxidised soybean oil

1.5 % of sodium butylnaphthalenesulfonate

2.5 % of sodium ligninsulfonate

42 % of terminally alkylated fatty alcohol alkoxyate (Plurafac[®] LF 431)

42 % of highly dispersed urea-formaldehyde polycondensate

Example A4 (formulation of a wettable powder)

35.0 % of compound of formula IV

1.1 % of sodium lauryl sulfate

3.0 % of sodium ligninsulfonate

20.9 % of precipitated silicic acid

40.0 % of isotridecanol 8 EO

The initial suspensibility is 70% and after storage for 6 months at room temperature is still 65 %.

Example A5 (formulation of a wettable powder)

30.0 g of compound of formula IV

1.1 g of sodium lauryl sulfate

3.0 g of sodium ligninsulfonate

- 12 -

22.5 g of precipitated silicic acid
1.9 g of pyrogenic silicic acid
40.0 g of isotridecanol 8 EO
1.5 g of calcined diatomaceous earth

The initial suspensibility is 90 % and after storage for 6 months at room temperature is still 85 %.

Example A6 (formulation of a wettable powder)

25.0 g of compound of formula IV
0.9 g of sodium lauryl sulfate
2.5 g of sodium ligninsulfonate
28.3 g of calcium silicate
1.6 g of pyrogenic silicic acid
40.0 g of isotridecanol 8 EO
1.7 g of calcined diatomaceous earth

The initial suspensibility is 74% and after storage for 6 months at room temperature is still 70 %.

The following Examples illustrate the biological activity of such dispersions.

B) Application Examples

Example B1

Postemergence herbicidal action

An aqueous dispersion of the formulation of Example A1 is sprayed in different concentrations, expressed in g of active ingredient per hectare, onto the plants in a greenhouse trial. The plants are then kept at 20°C and 45-60 % relative humidity and regularly watered. The trials are evaluated visually 23 days after treatment. The following species of grass are treated: *Avena fatua*, *Lolium perenne*. For control purposes, the wheat variety Arina is treated concurrently. The result is shown Table 1.

- 13 -

Table 1

Relative damage in % to untreated control plants
(visual assessment of the reduction of biomass)

	Concentration (g of active ingredient/ha)		
	40	20	10
"Arina"wheat	0	0	0
Alopecurus	95	90	30
Avena	95	75	0
Lolium	80	60	20

What is claimed is:

1. A formulation comprising

- a) at least one herbicide,
- b) at least one porous solid carrier material, and
- c) a nonionic surfactant,

which formulation contains 30 to 60 % by weight, based on the total formulation, of at least one nonionic surfactant of formula I



wherein R_1 is C_8 - C_{16} alkyl or C_8 - C_{16} alkenyl, R_2 is identical or different C_2 - C_4 alkylene, R_3 is hydrogen or C_1 - C_4 alkyl, and n is an integer from 5-10.

2. A formulation according to claim 1, wherein R_1 is C_{10} - C_{15} alkyl.

3. A formulation according to claim 1, wherein R_2 is a mixture of C_2 alkylene and C_4 alkylene, and R_3 is hydrogen or methyl.

4. A formulation according to claim 1, wherein R_2 is C_2 alkylene, R_3 is hydrogen, and n is an integer from 6 to 8.

5. A formulation according to claim 1, wherein the solid carrier material is selected from the group consisting of calcite, talcum, kaolin, montmorillonite, attapulgite, sepiolite, bentonite, highly dispersed silicic acid, synthetic highly absorbent silicic acid, and a highly dispersed urea/formaldehyde polycondensate.


6. A formulation according to claim 1, wherein the herbicide is selected from the group consisting of the group of the urea derivatives, imidazolinones, sulfonylureas, uracils, cyclohexanediones, chloroacetanilides, oxime ethers, triazines or phenoxypropionic acid derivatives.

7. A formulation according to claim 1, which additionally comprises a safener selected from the group of the quinoline derivatives.


8. A formulation according to claim 6, which comprises a herbicide selected from the group of the phenoxypropionic acid derivatives.

- 15 -

9. A formulation according to claim 1, wherein a compound of formula II is used in the form of the R enantiomer or the racemate



wherein Q is a radical of formula




wherein R_4 is hydrogen, F, Cl or Br;

R_5 is CF_3 , CN, Cl, Br or I;

R_7 and R_8 are each independently of the other F, Cl, Br, I, CF_3 or CN,

X is oxygen or sulfur, and n is 0 or an integer from 1 to 3;

Z is a radical of formula





wherein R_6 is hydrogen, C_1 - C_8 alkyl, C_3 - C_8 alkenyl, C_3 - C_8 alkynyl, C_1 - C_8 haloalkyl, C_3 - C_8 haloalkenyl, C_3 - C_8 haloalkynyl, or C_1 - C_6 alkyl which is substituted by CN, NO_2 , C_1 - C_4 alkoxy, C_1 - C_4 alkyl-SO-, C_1 - C_4 alkyl-SO₂-, C_1 - C_8 alkoxycarbonyl, C_3 - C_8 alkenyloxycarbonyl, C_3 - C_8 alkynyloxycarbonyl, C_1 - C_4 alkylidene-aminoxy or $N(R_{10})(R_{11})$, or is C_3 - C_8 alkenyl;


- 16 -

R₉ alkyl, haloalkyl, CN or C₁-C₄alkoxycarbonyl;
 and R₁₀ and R₁₁ are each independently of the other C₁-C₈alkyl, C₃-C₈alkenyl,
 C₃-C₈alkynyl, phenyl, benzyl or phenyl or benzyl, each substituted by halogen, NO₂, CN,
 OCH₃, or CF₃, or R₁₀ and R₁₁, together with the linking nitrogen atom, form a
 5-7-membered saturated heterocycle.

10. A composition according to claim 9, which comprises a compound of formula III or IV



11. A composition according to claim 10, which additionally comprises a compound of formula VIII



as safener.

12. A formulation according to claim 1, which contains the porous carrier material in an amount of 30 to 60 % by weight, based on the total formulation.

13. A formulation according to claim 1, which contains the porous carrier material in an amount of 35 to 50 % by weight, based on the total formulation.

- 17 -

14. A formulation according to claim 1, which contains the herbicide in an amount of 1 to 30 % by weight, based on the total formulation.
15. A formulation according to claim 1, which contains the herbicide in an amount of 5 to 15 % by weight, based on the total formulation.
16. A process for the preparation of a formulation as claimed in claim 1, which comprises mixing components a), b) and c) with one another.
17. A herbicidal composition, which contains a biologically active amount of the formulation as claimed in claim 1 dispersed in water.
18. A method of controlling weeds and grasses in crops of useful plants, which comprises treating said crops, the seeds or the crop area thereof with an effective amount of a herbicidal composition as claimed in claim 17.
19. Use of a formulation as claimed in claim 1 for the preparation of an aqueous, biologically active dispersion for selectively controlling weeds and grasses in crops of useful plants.
20. Use of a nonionic surfactant of formula I for the preparation of a wettable powder formulation for use as a herbicidal composition.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/IB 95/00001

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A01N25/30 A01N47/36 A01N43/60 A01N43/40 A01N43/42

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP,A,0 272 542 (BASF AG) 29 June 1988	1-19
X	see page 3, line 42-51 ---	20
Y	EP,A,0 573 898 (IDEMITSU KOSAN CO LTD) 15 December 1993	1-19
X	see page 8, line 2; example 5 ---	20
Y	US,A,5 171 352 (GROSSMANN K ET AL) 15 December 1992	1-19
X	see column 15, line 31 - column 16, line 25 ---	20
Y	EP,A,0 083 556 (CIBA GEIGY AG) 13 July 1983 cited in the application see page 9, paragraph 5 ---	8-10
	--- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

24 April 1995

Date of mailing of the international search report

24.05.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Klaver, J

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 95/00001

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP,A,0 191 736 (CIBA GEIGY AG) 20 August 1986 cited in the application see page 74, paragraph 4 ---	7-11
Y	EP,A,0 120 814 (CIBA GEIGY AG) 3 October 1984 cited in the application see p. 16, par. 5 and p. 18, par. 6 . -----	1-6, 12-19

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/IB 95/00001

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
EP-A-0272542	29-06-88	DE-A-	3643246	30-06-88
		AU-B-	600059	02-08-90
		AU-A-	8264887	23-06-88
		DE-A-	3784658	15-04-93
		ES-T-	2044903	16-01-94
		US-A-	4936899	26-06-90

EP-A-0573898	15-12-93	JP-A-	5339109	21-12-93
		AU-B-	649372	19-05-94
		US-A-	5348932	20-09-94

US-A-5171352	15-12-92	AU-A-	6015190	07-02-91
		CN-A-	1049439	27-02-91
		DE-A-	4023680	14-02-91
		ES-B-	2032252	16-01-94
		GR-B-	1000753	30-12-92
		IL-A-	95156	31-07-94
		TR-A-	24994	01-09-92

EP-A-0083556	13-07-83	CA-A-	1180717	08-01-85
		JP-C-	1502828	28-06-89
		JP-A-	58121273	19-07-83
		JP-B-	63052027	17-10-88
		US-A-	5002604	26-03-91
		US-A-	4505743	19-03-85
		US-A-	4713109	15-12-87
		US-A-	4935051	19-06-90

EP-A-0191736	20-08-86	AU-B-	600434	16-08-90
		AU-A-	5346686	21-08-86
		CZ-A-	8601008	16-11-94
		JP-A-	61191676	26-08-86
		RO-A-	100369	25-04-91
		US-A-	4881966	21-11-89

EP-A-0120814	03-10-84	AU-B-	570037	03-03-88
		AU-A-	2612284	04-10-84
		AU-B-	600331	09-08-90
		AU-A-	8311687	14-04-88
		CA-A-	1223589	30-06-87

information on patent family members

PCi/IB 95/00001

Form PCT/ISA/210 (patent family annex) (July 1992)

THIS PAGE BLANK (USPTO)